



What is bugging my fish? Some thoughts on managing fish pathogens

Department of Natural Resources
Gary E. Whelan



Why Care?



Annual
Mortality


=




+



Fish Pathogens and Disease



Disease Progression



Fish Pathogen “Rules”



- **Fish diseases issues are heavily regulated nationally and internationally**
 - Aquaculture driven
- **Water is a very good medium for transmission**
 - Constant movement of pathogens
 - Must manage acquired pathogens forever
- **Nearly none of the virus and bacteria are human pathogens**
 - Humans are too hot for replication of viruses and bacteria
 - Many parasites “can” use humans as intermediate hosts
- **Key pathogen groups require different strategies – Sampling and treatment**
 - Viruses
 - Bacteria
 - Parasites
 - Fungal
- **Make sure you really know you have the pathogen**
 - Confirm, confirm, confirm.....

Fish Pathogen “Rules”



- Poor knowledge base of baseline in wild
 - Understand disease ecology and progression
- Most wild epizootics are the result of ecosystem stress
- Fish stocking is one of the three legs of the management chair
 - Raise large numbers of fish – Disease management is critical
 - Herd Immunity
 - Use hatchery fish to break transmission
 - **Change natural mortality**
- Control vectors of transmission
- Fish pathogen expertise and control is organized
 - GLFC – Great Lakes Fish Health Committee
 - NE and NW Fish Health Committees

Fish Disease Transmission Relative Risk Factors



- Anglers

- Bait

- Preserved bait

- Very low to zero risk

- Frozen Bait

- Short and long distance
 - Low risk – Bacteria and some parasites
 - High risk – Viruses - depending on storage time

- Live Bait

- Short and long distance
 - Very high risk for all pathogens



Disease Transmission Relative Risk Factors



- Anglers

- Equipment

- Short and long distance
 - Depends on disinfection and pathogen

- Private and angler fish stocking

- Short and long distance
 - Depends on testing and biosecurity
 - Most private facilities are not a vector
 - Angler stocking is a large issue

- Bilge and live well water in recreational boats

- Short and long distance
 - Low risk but high frequency – Low amounts and needs perfect timing



Disease Transmission Relative Risk Factors



- Riparians

- Weed harvesting

- Short and long distance
 - Medium to High Risk depending on biosecurity

- Dredging

- Usually short distance
 - Depends on biosecurity





Disease Transmission Relative Risk Factors




- Commercial Shipping
 - Individual events have low probability but high frequency and volume activity with great potential effects
 - Short and long distance
 - Generally high volumes – 1 to 10 million gallons
 - Availability of enough viable pathogen at intake
 - Transportation viable
 - Available hosts at discharge
 - Dilution effects on enough viable pathogens at discharge

Disease Transmission Relative Risk Factors



- Resource Management Actions
 - Fish Stocking (State and Federal fish)
 - Short and long distance
 - Very low risk
 - Disinfect all eggs and inspect all fish
 - Movement of equipment
 - Short and long distance
 - Very low risk
 - Biosecurity




CO Whirling Disease Example

Disease Transmission Relative Risk Factors



- Fish migrations
 - Short and long distances possible
 - No stated evidence in literature – Sea Lice?
 - Generally low risk
- Fish eating birds and mammals
 - Depends on pathogen life cycle and residence time
 - Likely short distance
 - Risk - Variable
 - Low to Medium for viruses and bacteria
 - High for many parasites





Key Fish Pathogens



- Not all pathogens are equal
- Key wild pathogens
 - Viral Hemorrhagic Septicemia virus (VHS)
 - *Renibacterium salmoninarum* - Bacterial Kidney Disease
 - *Myxobolus cerebralis* (whirling disease)
 - Largemouth Bass Virus (LMBv)
 - Koi herpesvirus (KHV)
 - Spring Viraemia of Carp Virus (SVC)
- Key hatchery pathogens
 - *Aeromonids*
 - *Aeromonas salmonicida* (Furunculosis)
 - Flavobacterium
 - Infectious Salmon Anemia virus (ISA)
 - Infectious Haematopoietic Necrosis (IHN)

GLFHC Disease Transmission Risk Assessment Tool

| Factor | Score Range * Weight | Total Possible Score-650 |
|-----------------------------|---------------------------|--------------------------|
| Prevalence in Source | $(0 - 5) * 10$ | 50 |
| Transmission | $(1-5) * 10$ | 50 |
| Prevalence in Receiving | $(1-10) * 5$ | 50 |
| Regional Distribution | $(1-3) * 10$ | 30 |
| Number of Vectors | $(1-8+) * 10$ | 70 |
| Intro Increases Pathogen | $(0-5) * 5$ | 25 |
| Disease Spread to Others | $(0-5) * 10 + (0-5) * 10$ | 100 |
| Wild Stock Epizootic | $(1-5) * 20$ | 100 |
| Test Method Confidence | $(1-7) * 5$ | 35 |
| Health History of Source | $(1-5) * 5$ | 25 |
| Health History of Receiving | $(1-5) * 5$ | 25 |
| System Health History | $(1-5) * 5$ | 25 |
| Surveillance History | $(1-7) * 10$ | 70 |

GLFHC Disease Transmission Risk Assessment Tool

- Scoring and Recommendations



| Score | Risk | Recommendation |
|------------------|--------|--------------------------|
| Less Than 360 | Low | Unrestricted movement |
| 370 - 569 | Medium | Okay for positive waters |
| Greater Than 570 | High | No movement |






Questions or Comments

whelang@michigan.gov

Bacterial Kidney Disease Adventure Chronology



- 1955 – First reported in MI
- 1961 – BKD not likely to be spread by stocking
- 1967 – First mortalities of COS in Lake MI
 - Prevalence between 37.5 to 50% with dead salmon on beaches including CHS and Kokanee Salmon
- 1970 – Biosecurity measures started
 - Eliminating infected stocks, move away from open water sources



Bacterial Kidney Disease Adventure Chronology



- 1984 and 1985 – Small salmon noted at weirs
- 1986 and 1987 – Dead CHS in Lake MI
- 1988 – 10,000 to 20,000 CHS washed ashore in Lake MI with BKD detected (100% prevalence)
- 1989 – CHS mortality continues
 - BKD found in CHS smolts
 - Broodstock culling and “disease-free” eggs brought in from NY
 - **Mass Panic in Agency**

Bacterial Kidney Disease Adventure Chronology



- 1990-1991
 - Culling and water hardening in Erythromycin
 - CHS and COS broodstock BKD prevalence 100%
 - All production conducted with disease-free CHS eggs
 - Treating sick fish with Erythromycin feeds
- 1992-1994
 - WLSFH ATS develop BKD - 50,000K destroyed
 - Multiple phenotypes of BKD indentified
 - Reduced rations implicated as a stressor agent
 - Improved culling initiated using clinical signs and FELISA
 - Direct mortalities subside



Bacterial Kidney Disease

Adventure Chronology



- 1995-1999
 - Culling extended to all feral salmonid broodstocks
 - Salmon stocking reduced to 4.5 million
 - Salmon mortalities very low but in a broad range of hosts
- 2000-2002
 - BKD found in BKT from MSFH water source – 50K yearlings destroyed - \$50,000 loss (25%)
 - Changes
 - COS losses at PRSFH 12% to 4% (1992-1998) - \$160,000 savings
 - BKD dropped in CHS and COS from 15% to 3% (1991-1999)
 - BKD found in 51 of 67 streams sampled


Bacterial Kidney Disease Adventure Chronology



- 2002 to today – Increased biosecurity measures
 - Stress Reduction
 - Decreased rearing densities
 - Installed raceway covers
 - Improved water quality
 - Broodstock development
 - Pre-test donor populations
 - Testing individual pairs – culling most positive
 - Bringing new broodstocks to quarantine or isolation facilities
 - Increasing testing and vaccination of new broodstocks
 - Example of improvement - MSFH
 - BKT from >80% (>50% high) to 0% prevalence



Bacterial Kidney Disease Adventure Chronology



- 2002 to today – Increased biosecurity measures
 - Improved culling using “real time” ELISA
 - Increased testing during egg takes
 - Increased hygiene at hatcheries
 - Immediate removal of dead fish
 - Treatment of BKD positive fish
 - Increased cleaning and disinfection
 - UV disinfection
 - Water hardening of eggs in erythromycin

